Medications Quiz (3rd Year)

What is AF?		
What is the common anticoagulant use	ed in AF to minimise the risk	of clots?
What is the drug used to reverse the e	ffects of the above medication	on?
Are these types of insulin short, mediu	ım or long acting?	
Novorapid		
LevemirHumulin		
Lantus		
Which of these medications should not	t be given before theatre?	
Omeprazole	Clopidogrel	Levothyroxine
Bisoprolol	Zomorph	Domperidone
What is the reason for this?		
What is the difference between Phosp	hate Sandoz and Sando K?	

What should I do!?

you are caring for a patient with heart failure; they are bedematous and short of breath. Their blood pressure is 82/40. You look through their previous observations to find that this is sometimes normal for them. The patient is prescribed 80mg Furosemide . Do you give the medication? If yes or no, please give an explanation as to your decision.
One of your patients is prescribed Meropenem and they have a Penicillin allergy. What should you do?
You are caring for a patient who has come back from theatre at 16:00 following a DHS. You are administering medications at 18:00 and your patient is due Deltaparin . Before administering the mediation, what two things should you check?
You are working on nights and you are performing the controlled drug checks. When checking Oxycodone, you find that there should be 28 tablets. However, you can only find 14, therefore there is a strip missing. What actions should you take?
You are caring for a patient with type 1 diabetes and they are prescribed 10 units of Insulatard in the morning. You take their blood sugar before breakfast and it is 3.8. The patient states they do not feel like any breakfast this morning and may have something to eat later. Do you administer or omit the insulin?